# Recombinant Sialic Acid Binding Ig Like Lectin 10 (SIGLEC10) Instruction Manual

# SIPD328Hu01

## Homo sapiens (Human)

**Source** Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass20.1kDa

Accurate Molecular Mass 17kDa(Analysis of differences refer to the manual)

**Residues & Tags** Met571~Gln697 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits** Freeze-dried powder

Purity > 97% Isoelectric Point 9.5

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

## **SEQUENCE**

MKILPKRRTQ TETPRPRFSR HSTILDYINV VPTAGPLAQK RNQKATPNSP RTPLPPGAPS PESKKNQKKQ YQLPSFPEPK SSTQAPESQE SQEELHYATL NFPGVRPRPE ARMPKGTQAD YAEVKFQ

#### USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



# [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.