# Recombinant Tyrosine Aminotransferase (TAT) Instruction Manual

# SIPD348Hu01

### Homo sapiens (Human)

**Source** Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationCytoplasmPredicted Molecular Mass30.0kDa

Accurate Molecular Mass 33kDa(Analysis of differences refer to the manual)

**Residues & Tags** Cys221~Lys454 with N-terminal His Tag

100mMNaHCO<sub>3</sub>, 500mMNaCl, pH8.3, containing 1mM

**Buffer Formulation** EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and

Proclin300.

**Traits** Freeze-dried powder

Purity > 97% Isoelectric Point 5.5

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

### **SEQUENCE**

CGSVFSKRHL QKILAVAARQ CVPILADEIY

GDMVFSDCKY EPLATLSTDV PILSCGGLAK RWLVPGWRLG WILIHDRRDI FGNEIRDGLV KLSQRILGPC TIVQGALKSI LCRTPGEFYH NTLSFLKSNA DLCYGALAAI PGLRPVRPSG AMYLMVGIEM EHFPEFENDV EFTERLVAEQ SVHCLPATCF EYPNFIRVVI TVPEVMMLEA CSRIQEFCEQ HYHCAEGSQE

ECDK

### **USAGE**

Reconstitute in 100mM NaHCO3, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# **Image**



## [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.