Recombinant Thrombospondin 2 (THBS2) Instruction Manual

SIPD360Mu01

Mus musculus (Mouse)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass53.3kDa

Accurate Molecular Mass 53kDa(Analysis of differences refer to the manual)

Residues & Tags 53kDa(Analysis of differences refer to the manual)

Asp968~Arg1170 with N-terminal His and GST Tag

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 97% Isoelectric Point 6.4

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

Buffer Formulation

DFR NFQMVPLDPK GTTQIDPNWV IRHQGKELVQ TANSDPGIAV GFDEFGSVDF SGTFYVNTDR DDDYAGFVFG YQSSSRFYVV MWKQVTQTYW EDKPSRAYGY SGVSLKVVNS TTGTGEHLRN ALWHTGNTEG QVRTLWHDPK NIGWKDYTAY RWHLIHRPKT GYMRVLVHEG KQVMADSGPI YDQTYAGGRL GLFVFSQEMV YFSDLKYECR

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE


Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image

ENTIGENOUS RESERVATION ACTION TO THE PROPERTY OF CANDAGE CAMERICA SERVE OF CANDAGE CONTROL OF THE PROPERTY OF



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.