Recombinant Lipase A, Lysosomal Acid (LIPA) Instruction Manual

SIPD409Ra01

Rattus norvegicus (Rat)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Lysosome
Predicted Molecular Mass	41.1kDa
Accurate Molecular Mass	41kDa(Analysis of differences refer to the manual)
Residues & Tags	Asn32~Ile358 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 97%
Isoelectric Point	7.3
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

NMNVTEIIM HWGYPEHSVQ TGDGYILGVH RIPHGRKNQF DKGPKPVVYL QWRHGFLADS SNWVTNIDNN SLGFILADAG FDVWMGNSRG NTWSRKHKTL SVSQDEYWAF SFDEMAKYDL PASINYILNK TGQEQLYNVG HSQGCTIGFI AFSQMPELAK KVKMFFALAP VLSLNFASGP MVKLGRLPDL LLEDLFGQKQ FLPQSAMVKW LSTHICTHVI MKELCANIFF LICGFNEKNL NMSRVDVYTT HCPAGTSVQN MVHWTQVVKY HKLQAFDWGS SDKNYFHYNQ SYPPLYSIKD MQLPTALWSG GKDWLADTSD INILLTEI

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

STABILITY

Image

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.