

Recombinant Myosin VA (MYO5A) Instruction Manual

SIPD772Ra01

Rattus norvegicus (Rat)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	n/a
Predicted Molecular Mass	40.7kDa
Accurate Molecular Mass	41kDa(Analysis of differences refer to the manual)
Residues & Tags	Ser1504~Val1828 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	8.4
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
SLLTSTI NSIKKVLKKR GDDFETVSFW LSNTCRFLHC LKQYSGEEGF  
MKHNTRQNE HCLTNFDLAE YRQVLSDLAI QIYQQLVRVL ENILQPMIVS  
GMLEHETIQG VSGVKPTGLR KRTSSIADEG TYTLDSILRQ LNSFHSMVCQ  
HGMDPELIQ VVKQMFYIVG AITLNNLLLR KDMCSWSKGM QIRYNVSQLE  
EWLRDKNLMN SGAKETLEPL IQAAQLLQVK KKTDDDAEAI CSMCNALTTA  
QIVKVLNLYT PVNEFEERVS VSFIRTIQVR LRDRKDSPQL LMDAKHIFPV  
TFPFNPSSLA LETIQIPASL GLGFIARV
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.