

Recombinant N-Acylsphingosine Amidohydrolase 2 (ASAH2) Instruction Manual

SIPE261Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Membrane, Secreted, Golgi apparatus
Predicted Molecular Mass	50.4kDa
Accurate Molecular Mass	21&40&56kDa(Analysis of differences refer to the manual)
Residues & Tags	Ala451~Ile780 with Two N-terminal Tags, His-tag and SUMO-tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	6.7
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
ALGYSFAAGT IDGVGGLNFT QGKTEGDPFW DTIRDQILGK PSEEIKECHK  
PKPILLHTGE LSKPHPWHPD IVDVQIITLG SLAITAIPGE FTTMSGRRRLR  
EAVQAEFASH GMQNMTVVIS GLCNVYTHYI TTYEEYQAQR YEAASTIYGP  
HTLSAYIQLF RNLAKAIATD TVANLSRGPE PPFFKQLIVP LIPSIVDRAP  
KGRTFGDVLQ PAKPEYRVGE VAEVIFVGAN PKNSVQNQTH QTFLTVEKYE  
ATSTSWQIVC NDASWETRFY WHKGLLGLSN ATVEWHIPDT AQPGIYRIRY  
FGHNRKQDIL KPAVILSFEG TSPAFEVVTI
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.