Recombinant Eukaryotic Translation Initiation Factor 2 Alpha Kinase 3 (EIF2aK3) Instruction Manual

SIPE376Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass58.7kDa

Accurate Molecular Mass 59kDa(Analysis of differences refer to the manual)

Residues & Tags Phe593~Phe1077 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 95% Isoelectric Point 5.2

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

				FEPIQCLG
RGGFGVVFEA	KNKVDDCNYA	IKRIRLPNRE	LAREKVMREV	KALAKLEHPG
IVRYFNAWLE	APPEKWQEKM	DEIWLKDEST	DWPLSSPSPM	DAPSVKIRRM
DPFATKEHIE	IIAPSPQRSR	SFSVGISCDQ	TSSSESQFSP	LEFSGMDHED
ISESVDAAYN	LQDSCLTDCD	VEDGTMDGND	EGHSFELCPS	EASPYVRSRE
RTSSSIVFED	SGCDNASSKE	EPKTNRLHIG	NHCANKLTAF	KPTSSKSSSE
ATLSISPPRP	TTLSLDLTKN	TTEKLQPSSP	KVYLYIQMQL	CRKENLKDWM
NGRCTIEERE	RSVCLHIFLQ	IAEAVEFLHS	KGLMHRDLKP	SNIFFTMDDV
VKVGDFGLVT	AMDQDEEEQT	VLTPMPAYAR	HTGQVGTKLY	MSPEQIHGNS
YSHKVDIFSL	GLILFELLYP	FSTQMERVRT	LTDVRNLKFP	PLFTQKYPCE
YVMVQDMLSP	SPMERPEAIN	IIENAVF		

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.				