

Recombinant Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) Instruction Manual

SIPE934Ra01

Rattus norvegicus (Rat)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted, Lysosome, Cytoplasm, Golgi apparatus, Endoplasmic reticulum lumen, Endosome
Predicted Molecular Mass	58.1kDa
Accurate Molecular Mass	58kDa(Analysis of differences refer to the manual)
Residues & Tags	Trp164~Phe428 with N-terminal His and GST Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	5.9
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
WQQTEED SSPDGSSQVE VYLLDTSIQS GHREIEGRVT  
ITDFNSVPEE DGTRFHRQAS KCDSHGTHLA GVVSGRDAGV AKGTSLHSLR  
VLNCQGKGTV SGTLLIGLEFI RKSQLIQPSG PLVVLLPLAG GYSRILNTAC  
QRLARTGVVL VAAAGNFRDD ACLYSPASAP EVITVGATNA QDQPVTLGTL  
GTNFGRCVDL FAPGKDIIGA SSDCSTCYMS QSGTSQAAAH VAGIVAMMLN  
RDPALTLAEL RQRLILFSTK DVINMAWF
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

