

Recombinant 17-Beta-Hydroxysteroid Dehydrogenase Type 1 (HSD17b1) Instruction Manual

SIPF407Ra01

Rattus norvegicus (Rat)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Cytoplasm
Predicted Molecular Mass	50.7kDa
Accurate Molecular Mass	55kDa(Analysis of differences refer to the manual)
Residues & Tags	Met1~Ile344 with Two N-terminal Tags, His-tag and SUMO-tag
Buffer Formulation	PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	6.0
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
MDSTVVLITGCSSGIGLHLAVRLASDRSQSFVYATLRDLKSQGPLLEAARAQGCPPGSLEILELDVRDSESVAAARACVTEGRVDVL
VCNAGRGLFGPLEAHELNAVAVLDVNVLTGTIRMLQAFPLDMKRRHSGRVLVTASVGGMLMGLPFHEVYCASKFALEGLCESLAILLPL
FGVHVSLIECGAVHTAFHEKLEGGPGGALERADAQTRHLFAHYQRGYEQALSEAQDPEEVTELFLLTAMRAPQALRYFSTNRFLPLAR
MRTEDPSGSSVVEAMHREAFSDLQVQEGAKAGAQVSGDPDTPPRALICLPECAIPRVTAELGWSASDKPGQNKSCYQQKI
```

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.