

# Recombinant Brain Specific Angiogenesis Inhibitor 3 (BAI3) Instruction Manual

**SIPF711Hu01**

**Homo sapiens (Human)**

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Membrane
<b>Predicted Molecular Mass</b>	57.9kDa
<b>Accurate Molecular Mass</b>	58kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Cys30~Pro508 with N-terminal His Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 80%
<b>Isoelectric Point</b>	7.8
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

## SEQUENCE

C STLVKGVIYG SYSVSEMPK  
NFTNCTWLE NPDPTKYSIY LKFSKKDLSC SNFSLLAYQF DHFSHEKIKD  
LLRKNHSIMQ LCNSKNAFVF LQYDKNFIQI RRVFPTNFPG LQKKGEEDQK  
SFFEFLVLNK VSPSQFGCHV LCTWLESCLK SENGRTESCG IMYTKCTCPQ  
HLGEWGIDDQ SLILLNNVVL PLNEQTEGCL TQELQTTQVC NLTREAKRPP  
KEEFGMMGDH TIKSQRPRSV HEKRVPEQA DAAKFMAQTG ESGVEEWSQW  
STCSVTCGQG SQVRTRTCVS PYGTHCSGPL RESRVCNNTA LCPVHGVWEE  
WSPWSLCSFT CGRGQRTRTR SCTPPQYGGR PCEGPETHHK PCNIALCPVD  
GQWQEWSWS QCSVTCSNGT QQRSRQCTAA AHGGSECRGP WAESRECYNP  
ECTANGQWNQ WGHWSGCSSK CDGGWERRIR TCQGAIVITGQ QCEGTGEEVR  
RCNEQRCP

## USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


## **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **Image**



### 15% SDS-PAGE

SDS-PAGE Image

#### **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.