

Recombinant Mdm2 p53 Binding Protein Homolog (MDM2) Instruction Manual

SIPG176Mu01

Mus musculus (Mouse)

| | |
|---------------------------------|---|
| Source | Prokaryotic expression |
| Host | E.coli |
| Endotoxin Level | <1.0EU per 1µg (determined by the LAL method) |
| Subcellular Location | Nucleus, Cytoplasm |
| Predicted Molecular Mass | 38.2kDa |
| Accurate Molecular Mass | 48kDa(Analysis of differences refer to the manual) |
| Residues & Tags | Met1~Asn310 with N-terminal His Tag |
| Buffer Formulation | 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300. |
| Traits | Freeze-dried powder |
| Purity | > 95% |
| Isoelectric Point | 4.2 |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

SEQUENCE

```
MCNTNMSVST EGAASTSQIP ASEQETLVRP KPLLLKLLKS VGAQNPTYTM  
KEIIFYIGQY IMTKRLYDEK QQHIVYCSND LLGDVFGVPS FSVKEHRKIY  
AMIYRNLVAV SQQDSGTSL SRRQPEGGS DLKDPLQAPP EEKPSSSDLI  
SRLSTSSRRR SISETEENTD ELPGERHRKR RRSLSFDPSL GLCELREMCS  
GGSSSSSSSS SESETETPSHQ DLDDGVSEHS GDCLDQDSVS DQFSVEFEVE  
SLDSEDYSL SDEGHELSD EDDEVYRVTVYQ TGESDTDSFE GDPEISLADY  
WKCTSCNEMN
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.