# Recombinant ATPase, H+ Transporting, Lysosomal Accessory Protein 2 (ATP6AP2) Instruction Manual

# SIPG416Hu01

### Homo sapiens (Human)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass35.4kDa

Accurate Molecular Mass 35kDa(Analysis of differences refer to the manual)

**Residues & Tags** Asn17~Glu302 with N-terminal His Tag

Buffer Formulation PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5%

Trehalose and Proclin300.

**Traits** Freeze-dried powder

Purity > 90% Isoelectric Point 6.4

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

## **SEQUENCE**

NEFS ILKSPGSVVF RNGNWPIPGE RIPDVAALSM
GFSVKEDLSW PGLAVGNLFH RPRATVMVMV KGVNKLALPP GSVISYPLEN
AVPFSLDSVA NSIHSLFSEE TPVVLQLAPS EERVYMVGKA NSVFEDLSVT
LRQLRNRLFQ ENSVLSSLPL NSLSRNNEVD LLFLSELQVL HDISSLLSRH
KHLAKDHSPD LYSLELAGLD EIGKRYGEDS EQFRDASKIL VDALQKFADD
MYSLYGGNAV VELVTVKSFD TSLIRKTRTI LEAKQAKNPA SPYNLAYKYN
FE

### **USAGE**


Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



# [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.