Recombinant Corin (CRN) Instruction Manual

SIPG734Hu03

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Membrane, Secreted

Predicted Molecular Mass 53.1kDa

Accurate Molecular Mass 53kDa(Analysis of differences refer to the manual)

Residues & Tags Arg134~Pro573 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 4.8

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

			RNTSACM	NITHSQCQML
PYHATLTPLL	SVVRNMEMEK	FLKFFTYLHR	LSCYQHIMLF	GCTLAFPECI
IDGDDSHGLL	PCRSFCEAAK	EGCESVLGMV	NYSWPDFLRC	SQFRNQTESS
NVSRICFSPQ	QENGKQLLCG	RGENFLCASG	ICIPGKLQCN	GYNDCDDWSD
EAHCNCSENL	FHCHTGKCLN	YSLVCDGYDD	CGDLSDEQNC	DCNPTTEHRC
GDGRCIAMEW	VCDGDHDCVD	KSDEVNCSCH	SQGLVECRNG	QCIPSTFQCD
GDEDCKDGSD	EENCSVIQTS	CQEGDQRCLY	NPCLDSCGGS	SLCDPNNSLN
NCSQCEPITL	ELCMNLPYNS	TSYPNYFGHR	TQKEASISWE	SSLFPALVQT
NCYKYLMFFS	CTILVPKCDV	NTGEHIPPCR	ALCEHSKERC	ESVLGIVGLQ
WPEDTDCSQF	PEENSDNQTC	LMP		

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.






Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.