Recombinant Trans Golgi Network Protein 2 (TGOLN2) Instruction Manual

SIPH019Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass34.7kDa

Accurate Molecular Mass 56kDa(Analysis of differences refer to the manual)

Residues & Tags Ala22~Glu323 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 97% Isoelectric Point 4.8

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

ATESVKQEE AGVRPSAGNV STHPSLSQRP
GGSTKSHPEP QTPKDSPSKS SAEAQTPEDT PNKSGAEAKT QKDSSNKSGA
EAKTQKGSTS KSGSEAQTTK DSTSKSHPEL QTPKDSTGKS GAEAQTPEDS
PNRSGAEAKT QKDSPSKSGS EAQTTKDVPN KSGADGQTPK DGSSKSGAED
QTPKDVPNKS GAEKQTPKDG SNKSGAEEQG PIDGPSKSGA EEQTSKDSPN
KEEVKSSEPT EDVEPKEAED DDTGPEEGSP PKEEKEKMSG SASSENREGT
LSDSTGSEKD DLYPNGSGNG SAE

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE


Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.







SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.