Recombinant Microtubule Associated Serine/Threonine Kinase 2 (MAST2) Instruction Manual

SIPH680Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Membrane, Cytoplasm

Predicted Molecular Mass 61.2kDa

Accurate Molecular Mass

Residues & Tags

Buffer Formulation

61kDa(Analysis of differences refer to the manual)

Phe512~Phe785 with N-terminal His and GST Tag

PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 5.4

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

FETIKLISN GAYGAVFLVR HKSTRQRFAM KKINKQNLIL RNQIQQAFVE RDILTFAENP FVVSMFCSFD TKRHLCMVME YVEGGDCATL LKNIGALPVD MVRLYFAETV LALEYLHNYG IVHRDLKPDN LLITSMGHIK LTDFGLSKIG LMSLTTNLYE GHIEKDAREF LDKQVCGTPE YIAPEVILRQ GYGKPVDWWA MGIILYEFLV GCVPFFGDTP EELFGQVISD EIVWPEGDEA LPPDAQDLTS KLLHONPLER LGTGSAYEVK QHPFF

USAGE

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.