Recombinant Inter Alpha-Globulin Inhibitor H1 (ITIH1) Instruction Manual

SIPH886Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass38.5kDa

Accurate Molecular Mass 38kDa(Analysis of differences refer to the manual)

Residues & Tags Ser507~Asp819 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 5.4

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
SEIV VAGRIADNKQ SSFKADVQAH GEGQEFSITC LVDEEMKKL
LRERGHMLEN HVERLWAYLT IQELLAKRMK VDREERANLS SQALQMSLDY
GFVTPLTSMS IRGMADQDGL KPTIDKPSED SPPLEMLGPR RTFVLSALQP
SPTHSSSNTQ RLPDRVTGVD TDPHFIIHVP QKEDTLCFNI NEEPGVILSL
VQDPNTGFSV NGQLIGNKAR SPGQHDGTYF GRLGIANPAT DFQLEVTPQN
ITLNPGFGGP VFSWRDQAVL RQDGVVVTIN KKRNLVVSVD DGGTFEVVLH
RVWKGSSVHO DFLGFYVLD
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.