Recombinant Plastin 1 (PLS1) Instruction Manual

SIPH955Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationCytoplasmPredicted Molecular Mass33.8kDa

Accurate Molecular Mass 34kDa(Analysis of differences refer to the manual)

Residues & Tags Thr108~Lys380 with N-terminal His Tag

Buffer Formulation PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5%

Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 97% Isoelectric Point 4.7

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

TST ISSEGTQHSY SEEKVAFVN WINKALENDP DCKHLIPMNP
NDDSLFKSLA DGILLCKMIN LSEPDTIDER AINKKKLTPF TISENLNLAL
NSASAIGCTV VNIGASDLKE GKPHLVLGLL WQIIKVGLFA DIEISRNEAL
IALLNEGEEL EELMKLSPEE LLLRWVNYHL TNAGWHTISN FSQDIKDSRA
YFHLLNQIAP KGGEDGPAIA IDLSGINETN DLKRAGLMLQ EADKLGCKQF
VTPADVVSGN PKLNLAFVAN LFNTYPCLHK

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.






Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.