

# Recombinant Nicotinamide-N-Methyltransferase (NNMT) Instruction Manual

**SIPH968Hu01**

**Homo sapiens (Human)**

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Cytoplasm
<b>Predicted Molecular Mass</b>	58.8kDa
<b>Accurate Molecular Mass</b>	59kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Met1~Arg258 with N-terminal His and GST Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 95%
<b>Isoelectric Point</b>	5.0
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

## SEQUENCE

```
MESGFTSKDT YLSHFNPRDY LEKYYKFGSR HSAESQILKH LLKNLFKIFC  
LDGVKGDLLI DIGSGPTIYQ LLSACESFKE IVVDYSDQN LQELEKWLKK  
EPEAFDWSPV VTYVCDLEGN RVKGPEKEEK LRQAVKQVLK CDVTQSQPLG  
AVPLPPADCV LSTLCLDAAC PDLPTYCRAL RNLGSLKPG GFLVIMDALK  
SSYYMIGEQQ FSSLPLGREA VEA AVKEAGY TIEWFEVISQ SYSSTMANNE  
GLFSLVAR
```

## USAGE

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.7 mg/mL. Do not vortex.

## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image




Figure. SDS-PAGE

## **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.