Recombinant Periostin (POSTN) Instruction Manual

SIPH997Hu03

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass47.3kDa

Accurate Molecular Mass 47kDa(Analysis of differences refer to the manual)

Residues & Tags Gly234~Leu628 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 5.0

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

GTSIQDF IEAEDDLSSF
RAAAITSDIL EALGRDGHFT LFAPTNEAFE KLPRGVLERI MGDKVASEAL
MKYHILNTLQ CSESIMGGAV FETLEGNTIE IGCDGDSITV NGIKMVNKKD
IVTNNGVIHL IDQVLIPDSA KQVIELAGKQ QTTFTDLVAQ LGLASALRPD
GEYTLLAPVN NAFSDDTLSM DQRLLKLILQ NHILKVKVGL NELYNGQILE
TIGGKQLRVF VYRTAVCIEN SCMEKGSKQG RNGAIHIFRE IIKPAEKSLH
EKLKQDKRFS TFLSLLEAAD LKELLTQPGD WTLFVPTNDA FKGMTSEEKE
ILIRDKNALQ NIILYHLTPG VFIGKGFEPG VTNILKTTQG SKIFLKEVND
TLLVNELKSK ESDIMTTNGV IHVVDKLL

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.