Recombinant Follistatin Like Protein 3 (FSTL3) Instruction Manual

SIPJ055Mu01

Mus musculus (Mouse)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Nucleus, Secreted

Predicted Molecular Mass 28.6kDa

Accurate Molecular Mass 29kDa(Analysis of differences refer to the manual)

Residues & Tags Val24~Val256 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 95% Isoelectric Point 7.8

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

VMGSEDS VPGGVCWLQQ GREATCSLVL

KTRVSREECC ASGNINTAWS NFTHPGNKIS LLGFLGLVHC LPCKDSCDGV ECGPGKACRM LGGRPHCECV PNCEGLPAGF QVCGSDGATY RDECELRTAR CRGHPDLRVM YRGRCQKSCA QVVCPRPQSC LVDQTGSAHC VVCRAAPCPV PSNPGQELCG NNNVTYISSC HLRQATCFLG RSIGVRHPGI CTGGPKVPAE

EEENFV

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.