Recombinant GLI Family Zinc Finger Protein 3 (GLI3) Instruction Manual

SIPJ454Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Nucleus, Cytoplasm

Predicted Molecular Mass 29.3kDa

Accurate Molecular Mass 29kDa(Analysis of differences refer to the manual)

Residues & Tags Met111~Ser343 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 8.9

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

	MDPRNGYMEP	HYHPPHLFPA	FHPPVPIDAR	HHEGRYHYDP
SPIPPLHMTS	ALSSSPTYPD	LPFIRISPHR	NPTAASESPF	SPPHPYINPY
MDYIRSLHSS	PSLSMISATR	GLSPTDAPHA	GVSPAEYYHQ	MALLTGQRSP
YADIIPSAAT	AGTGAIHMEY	LHAMDSTRFS	SPRLSARPSR	KRTLSISPLS
DHSFDLQTMI	RTSPNSLVTI	LNNSRSSSSA	SGSYGHLSAS	AIS

USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.