Recombinant Enhancer Of Zeste Homolog 2 (EZH2) Instruction Manual

SIPK047Hu02

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationNucleusPredicted Molecular Mass29.2kDa

Accurate Molecular Mass 33kDa(Analysis of differences refer to the manual)

Residues & Tags Arg494~Val737 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL,

5% Trehalose.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 8.3

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

				RKHRLWA
AHCRKIQLKK	DGSSNHVYNY	QPCDHPRQPC	DSSCPCVIAQ	NFCEKFCQCS
SECQNRFPGC	RCKAQCNTKQ	CPCYLAVREC	DPDLCLTCGA	ADHWDSKNVS
CKNCSIQRGS	KKHLLLAPSD	VAGWGIFIKD	PVQKNEFISE	YCGEIISQDE
ADRRGKVYDK	YMCSFLFNLN	NDFVVDATRK	GNKIRFANHS	VNPNCYAKVM
MVNGDHRIGI	FAKRAIQTGE	ELFFDYRYSQ	ADALKYV	

USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.