

Recombinant SPARC Related Modular Calcium Binding Protein 1 (SMOC1) Instruction Manual

SIPN970Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	75kDa
Accurate Molecular Mass	83kDa(Analysis of differences refer to the manual)
Residues & Tags	His27~Val434 with N-terminal His and GST Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	8.0
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
                HRTT GPRFLISDRD PQC�NLHCSRT
QPKPICASDG RSYESMCEYQ RAKCRDPTLG VVHRGRCKDA GQSKCRLERA
QALEQAKKPQ EAVFVPECGE DGSFTQVQCH TYTGWCWCVT PDGKPISGSS
VQNKTPVCSG SVTDKPLSQG NSGRKDDGSK PTPTMETQPV FDGDEITAPT
LWIKHLVIKD SKLNNTNIRN SEKVYSCDQE RQSALEEAQQ NPREGIVIPE
CAPGGLYKPV QCHQSTGYCW CVLVDTGRPL PGTSTRYVMP SCESDARAKT
TEADDPFKDR ELPGCPEGKK MEFITSLLLDA LTTDMVQAIN SAAPTGGGRF
SEPDPSHTLE ERVVHWYFSQ LDSNSSNDIN KREMKPFKRY VKKKAKPKKC
ARRFTDYCDL NKDKVISLPE LKGCLGVSKE GRLV
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.