Recombinant RAD51 Homolog (RAD51) Instruction Manual

SIPP878Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Nucleus, Mitochondrion, Cytoplasm

Predicted Molecular Mass 40.5kDa

Accurate Molecular Mass 41kDa(Analysis of differences refer to the manual)

Residues & Tags Ala2~Asp339 with N-terminal His Tag

Buffer Formulation PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Traits Freeze-dried powder

> 97% **Purity** 5.4 **Isoelectric Point**

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

AMOMOLEAN ADTSVEEESF GPOPISRLEO CGINANDVKK LEEAGFHTVE AVAYAPKKEL INIKGISEAK ADKILAEAAK LVPMGFTTAT EFHQRRSEII QITTGSKELD KLLQGGIETG SITEMFGEFR TGKTQICHTL AVTCQLPIDR GGGEGKAMYI DTEGTFRPER LLAVAERYGL SGSDVLDNVA YARAFNTDHQ TOLLYOASAM MVESRYALLI VDSATALYRT DYSGRGELSA ROMHLARFLR MLLRLADEFG VAVVITNOVV AQVDGAAMFA ADPKKPIGGN IIAHASTTRL

YLRKGRGETR ICKIYDSPCL PEAEAMFAIN ADGVGDAKD

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image

CATION CAST TAKEN TO SEAT TAKEN AGAIC TIGOTOMICATI ACTION GOTOMICATION CONTACT CONTACT




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.