

Recombinant Histone Cluster 1, H2ac (HIST1H2AC) Instruction Manual

SIPQ490Hu01

Homo sapiens (Human)

| | |
|---------------------------------|---|
| Source | Prokaryotic expression |
| Host | E.coli |
| Endotoxin Level | <1.0EU per 1µg (determined by the LAL method) |
| Subcellular Location | n/a |
| Predicted Molecular Mass | 41.0kDa |
| Accurate Molecular Mass | 44kDa(Analysis of differences refer to the manual) |
| Residues & Tags | Ser2~Lys130 (Accession # Q93077) with N-terminal His and GST Tag |
| Buffer Formulation | PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. |
| Traits | Freeze-dried powder |
| Purity | > 95% |
| Isoelectric Point | n/a |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

USAGE

Reconstitute in PBS or others.

STORAGE


Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.