Recombinant Isthmin 1 (ISM1) Instruction Manual

SIPQ569Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass79kDa

Accurate Molecular Mass 89kDa(Analysis of differences refer to the manual)
Residues & Tags Ala30~Tyr464 with N-terminal His and GST Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 4.9

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

A DGPDAAAGNA SQAQLQNNLN
VGSDTTSETS FSLSKEAPRE HLDHQAAHQP FPRPRFRQET GHPSLQRDFP
RSFLLDLPNF PDLSKADING QNPNIQVTIE VVDGPDSEAD KDQHPENKPS
WSVPSPDWRA WWQRSLSLAR ANSGDQDYKY DSTSDDSNFL NPPRGWDHTA
PGHRTFETKD QPEYDSTDGE GDWSLWSVCS VTCGNGNQKR TRSCGYACTA
TESRTCDRPN CPGIEDTFRT AATEVSLLAG SEEFNATKLF EVDTDSCERW
MSCKSEFLKK YMHKVMNDLP SCPCSYPTEV AYSTADIFDR IKRKDFRWKD
ASGPKEKLEI YKPTARYCIR SMLSLESTTL AAQHCCYGDN MQLITRGKGA
GTPNLISTEF SAELHYKVDV LPWIICKGDW SRYNEARPPN NGQKCTESPS
DEDYIKOFQE AREY

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image

WITH STREET THE RESIDENCE AND DESCRIPTION OF THE ANGEL AND A TREET AS AND A THE FOR THE ANGEL AND A SANDER THE FOR THE ANGEL AND A SANDE A




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.