

# Recombinant Activating Signal Cointegrator 1 Complex Subunit 3 (ASCC3) Instruction Manual

**SIPQ860Hu01**

**Homo sapiens (Human)**

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Nucleus, Cytoplasm
<b>Predicted Molecular Mass</b>	45.5kDa
<b>Accurate Molecular Mass</b>	46kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Pro1812~Tyr2176 with N-terminal His Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 95%
<b>Isoelectric Point</b>	7.1
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

## SEQUENCE

```
          PLTYGRIAS  YYYLKHQTVK  MFKDRLKPEC  STEELLSILS
DAEEYTDLPV  RHNE DHMNSE  LAKCLPIESN  PHSFDSPTK  AHLLLQAHLS
RAMLPDYPDY  TDTKT VLDQA  LRVCQAMLDV  AANQGWLTV  LNITNLIQMV
IQGRWLKDSS  LLTLPNIENH  HLHLFKKWKP  IMKGP HARGR  TSIESLPELI
HACGGKDHFV  SSMVESELHA  AKTKQAWNFL  SHLPVINVGI  SVKGSWDDL
EGHNELSVST  LTADKRDDNK  WIKLHADQEY  VLQVSLQRVH  FGFHKGKPES
CAVTPRFPKS  KDEGWFLILG  EVDKRELIAL  KRVG YIRNHH  VASLSFYTPE
IPGRYIYTLY  FMSDCYLGLD  QQYDIY
```

## USAGE

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image




Figure. SDS-PAGE

## **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.