

# Recombinant Sialic Acid Binding Ig Like Lectin 15 (SIGLEC15) Instruction Manual

**SIPR167Hu01**

**Homo sapiens (Human)**

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Membrane
<b>Predicted Molecular Mass</b>	56.8kDa
<b>Accurate Molecular Mass</b>	60kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Phe20~Leu266 with N-terminal His and GST Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 90%
<b>Isoelectric Point</b>	8.4
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

## SEQUENCE

```
F VRTKIDTTEN LLNTEVHSSP AQRWSMQVPP  
EVSAEAGDAA VLPCTFTHPH RHYDGPLTAI WRAGEPYAGP QVFRCAAARG  
SELCQTALSL HGRFRLLGNP RRNDLSLRVE RLALADDRRY FCRVEFAGDV  
HDRYESRHGV RLHVTAAPRI VNISVLPSPA HAFRALCTAE GEPPPALAWS  
GPALGNSLAA VRSPREGHGH LVTAELPALT HDGRYTCTAA NSLGRSEASV  
YLFRFHGASG ASTVAL
```

## USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.  
Do not vortex.

## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image

G A T T C G G A T C C C C G A T A T C C C A T G G A C C C G C G T C C C T C A A T A C C G G A G C C A C C C C G G T A C C C A G A T C T G G G C




Figure. SDS-PAGE

## **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

