

# Recombinant Cancerous Inhibitor Of PP2A (CIP2A) Instruction Manual

**SIPR567Hu02**

**Homo sapiens (Human)**

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Membrane, Cytoplasm
<b>Predicted Molecular Mass</b>	57.2kDa
<b>Accurate Molecular Mass</b>	50kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Asn230~Glu560 with His and TRxA Tag
<b>Buffer Formulation</b>	PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 90%
<b>Isoelectric Point</b>	5.0
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

## SEQUENCE

```

                                     N IHQTFQLIFN ILINGDGLT
RKYSVDLLMD LLKNPKIADY LTRYEHFSSC LHQVLGLLNG KDPDSSSKVL
ELLLAFCSVT QLRHMLTQMM FEQSPPGSAT LGSHTKCLEP TVALLRWLSQ
PLDGSENCVS LALELFKEIF EDVIDAANCS SADRFVTLLE PTILDQLQFT
EQNLDEALTR KKCERIAKAI EVLLTLCGDD TLKMHIKIL TTVKCTTLE
QQFTYQKIDL GFGTKVADSE LCKLAADVIL KTLDLINKLK PLVPGMEVSF
YKILQDPRLI TPLAFALTSD NREQVQSLR ILLEAAPLPD FPALVLGESI
AANNAYRQQE
```

## USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image




Figure. SDS-PAGE

## **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.