Recombinant Hyaluronoglucosaminidase 1 (HYAL1) Instruction Manual

SIPS062Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass47.7kDa

Accurate Molecular Mass 43kDa(Analysis of differences refer to the manual)

Residues & Tags Phe22~Trp435 with N-terminal His Tag

Buffer Formulation PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5%

Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 97% Isoelectric Point 6.5

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

		FRGPLLPNR	PFTTVWNANT	QWCLERHGVD
VDVSVFDVVA	NPGQTFRGPD	MTIFYSSQLG	TYPYYTPTGE	PVFGGLPQNA
SLIAHLARTF	QDILAAIPAP	DFSGLAVIDW	EAWRPRWAFN	WDTKDIYRQR
SRALVQAQHP	DWPAPQVEAV	AQDQFQGAAR	AWMAGTLQLG	RALRPRGLWG
FYGFPDCYNY	DFLSPNYTGQ	CPSGIRAQND	QLGWLWGQSR	ALYPSIYMPA
VLEGTGKSQM	YVQHRVAEAF	RVAVAAGDPN	LPVLPYVQIF	YDTTNHFLPL
DELEHSLGES	AAQGAAGVVL	WVSWENTRTK	ESCQAIKEYM	DTTLGPFILN
VTSGALLCSQ	ALCSGHGRCV	RRTSHPKALL	LLNPASFSIQ	LTPGGGPLSL
RGALSLEDOA	OMAVEFKCRC	YPGWOAPWCE	RKSMW	

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.