

# Recombinant Mab21 Domain Containing Protein 1 (MB21D1) Instruction Manual

**SIPU035Mu01**

**Mus musculus (Mouse)**

|                                 |   |
|---------------------------------|---|
| <b>Source</b>                   | Prokaryotic expression  |
| <b>Host</b>                     | E.coli  |
| <b>Endotoxin Level</b>          | <1.0EU per 1µg (determined by the LAL method)   |
| <b>Subcellular Location</b>     | Cytoplasm   |
| <b>Predicted Molecular Mass</b> | 24.9kDa   |
| <b>Accurate Molecular Mass</b>  | 27kDa(Analysis of differences refer to the manual)  |
| <b>Residues &amp; Tags</b>      | Glu211~Lys395 with N-terminal His Tag   |
| <b>Buffer Formulation</b>       | 100mMNaHCO <sub>3</sub> , 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300. |
| <b>Traits</b>                   | Freeze-dried powder   |
| <b>Purity</b>                   | > 95%   |
| <b>Isoelectric Point</b>        | 8.8   |
| <b>Applications</b>             | Positive Control; Immunogen; SDS-PAGE; WB.  |

## SEQUENCE

```
EFDVMFKLEV PRIELQEYYE TGAFYLVKFK RIPRGNPLSH  
FLEGEVLSAT KMLSKFRKII KEEVKEIKDI DVSVEKEKPG SPAVTLLIRN  
PEEISVDIIL ALESKGSWPI STKEGLPIQG WLGTKVRTNL RREPFYLVPK  
NAKDGNSFQG ETWRLSFSHT EKYILNNHGI EKTCESSGA KCCRK
```

## USAGE

Reconstitute in 100mM NaHCO<sub>3</sub>, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **Image**



15% SDS-PAGE

SDS-PAGE Image

## **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.